

8508 Straight Master

A straight is a poker hand containing five cards of sequential rank, not necessarily to be the same suit. For example, a hand containing 7 club, 6 spade, 5 spade, 4 heart and 3 diamond forms a straight. In this problem, we extend the definition of a straight to allow 3 to 5 cards of sequential rank. Hence a hand containing 'K' spade, 'Q' club, and 'J' heart is also a straight.

Mr. Panda is playing a poker game called Straight Master. The game uses a large deck of card that has N ranks from 1 to N . The rule of the game is simple: split the cards in Mr. Panda's hand into several straights of length from 3 to 5.

Now given a hand of cards, can you help Mr. Panda to determine if it is possible to split the cards into straights?

Input

The first line of the input gives the number of test cases, T . T test cases follow.

Each test case contains two lines. The first line contains an integer N , indicating the number of ranks in the deck. The next line contains N integers a_1, a_2, \dots, a_N indicating the number of cards for each rank in Mr. Panda's hand.

Output

For each test case, output one line containing 'Case # x : y ', where x is the test case number (starting from 1) and y is 'Yes' if Mr. Panda can split all his cards into straights of length from 3 to 5, or 'No' otherwise.

Limits:

- $1 \leq T \leq 100$.
- $1 \leq N \leq 2 \times 10^5$.
- $0 \leq a_i \leq 10^9$.
- $\sum N \leq 4 \times 10^6$.

Note

In the first test case, Mr. Panda can split his cards into two straights: [1, 2, 3] and [2, 3, 4].

In the second test case, there is no way to form a straight for card 6 and 7.

Sample Input

```
2
13
1 2 2 1 0 0 0 0 0 0 0 0 0
13
1 1 1 1 0 1 1 0 0 0 0 0 0
```

Sample Output

```
Case #1: Yes
Case #2: No
```